

Hadron and Nuclear Physics

Exercise sheet 1

Winter term 2013/14

Hand-in: In the exercise group on Nov 5th 2013

Problem 1.1: Decay of the Δ -Resonance (12 points)

- (a) In pion-nucleon scattering at a center-of-mass energy of 1232 MeV, the short-lived Δ -resonance is formed which was discovered in 1951 by Fermi and coworkers. The Δ^{++} decays according to

$$\Delta^{++} \rightarrow p + \pi^+,$$

where $m_\pi = 139.570 \text{ MeV}/c^2$.

Calculate, by applying relativistic kinematics to the decay of the Δ -resonance at rest, the energies and momenta of the final state particles as well as their kinetic energies and "velocities" β_p and β_π .

Problem 1.2: Conservation Laws (12 points)

- (a) Examine the following processes, and state for each one whether it is *possible* or *impossible*, according to the Standard Model. In the former case, state which interaction is responsible - strong, electromagnetic, or weak; in the latter case, cite a conservation law that prevents it from occurring.

a) $p + \bar{p} \rightarrow \pi^+ + \pi^0$

b) $\eta \rightarrow \gamma + \gamma$

c) $\Sigma^0 \rightarrow \Lambda + \pi^0$

d) $\Sigma^- \rightarrow n + \pi^-$

e) $e^+ + e^- \rightarrow \mu^+ + \mu^-$

f) $\mu^- \rightarrow e^- + \bar{\nu}_e$

g) $\Delta^+ \rightarrow p + \pi^0$

h) $\bar{\nu}_e + p \rightarrow n + e^+$

i) $e^- + p \rightarrow \nu_e + \pi^0$

j) $p + p \rightarrow \Sigma^+ + n + K^0 + \pi^+ + \pi^0$

k) $p \rightarrow e^+ + \gamma$

l) $p + p \rightarrow p + p + p + \bar{p}$

m) $n + \bar{n} \rightarrow \pi^+ + \pi^- + \pi^0$

n) $\pi^+ + n \rightarrow \pi^- + p$

o) $K^- \rightarrow \pi^- + \pi^0$

p) $\Sigma^+ + n \rightarrow \Sigma^- + p$

q) $\Sigma^0 \rightarrow \Lambda + \gamma$

r) $\Xi^- \rightarrow \Lambda + \pi^-$

s) $\Xi^0 \rightarrow p + \pi^-$

t) $\pi^- + p \rightarrow \Lambda + K^0$

u) $\pi^0 \rightarrow \gamma\gamma$

v) $\Sigma^- \rightarrow n + e^- + \bar{\nu}_e$

w) $n \rightarrow p + \mu^- + \bar{\nu}_\mu$

x) $\Delta^+ \rightarrow p$

Problem 1.3: Isospin (12 points)

- (a) Find the ratio of the cross sections of the following reactions, assuming the CM energy is such that $I = \frac{3}{2}$ dominates:

a) $\pi^- + p \rightarrow K^0 + \Sigma^0$


b) $\pi^- + p \rightarrow K^+ + \Sigma^-$

c) $\pi^+ + p \rightarrow K^+ + \Sigma^+$.

What if the energy is such that the $I = \frac{1}{2}$ channel dominates?

1 Übung: Besprechung

1.1 Aufgabe




$$\begin{aligned} \hbar = c = 1 \\ p_p = (E_p, p, 0, 0) \\ s = (p_p + p_\pi)^2 = (E_p + E_\pi)^2 \Rightarrow \sqrt{s} = E_p + E_\pi \\ p_p^2 = m_p^2 = E_p^2 - p^2, \quad p_\pi^2 = m_\pi^2 = E_\pi^2 - p^2 \end{aligned}$$

Impulserhaltung:

$$\begin{aligned} p^2 = p^2 \Leftrightarrow E_p^2 - m_p^2 = E_\pi^2 - m_\pi^2 \\ \Leftrightarrow E_p^2 - m_p^2 = E_\pi^2 - m_\pi^2 = (E_p - m_p)(E_p + m_p) = \sqrt{s}(E_p - m_p) \\ E_p = \frac{1}{2}(\sqrt{s} + \frac{m_p^2 - m_\pi^2}{\sqrt{s}}) \\ E_\pi = \frac{1}{2}(\sqrt{s} + \frac{m_\pi^2 - m_p^2}{\sqrt{s}}) \\ T_p = E_p - m_p, \quad T_\pi = E_\pi - m_\pi \\ p = \sqrt{E_p^2 - m_p^2} = p_\pi \\ \beta_p = \frac{p}{E_p}, \quad \beta_\pi = \frac{p}{E_\pi} \end{aligned}$$

1.2 Aufgabe: Teile




e) $e^+ + e^- \rightarrow \mu^+ + \mu^-$:

j) $p + p \rightarrow \Sigma^+ + n + K^0 + \pi^+ + \pi^0$: stark, Parität: je 2 Teilchen können 1 beitragen \Rightarrow erhalten.

1.3 Aufgabe: Besprechung

Toy-Modell:



$$\begin{pmatrix} p \\ n \end{pmatrix} \quad \begin{pmatrix} K^+ \\ K^0 \end{pmatrix} \quad \begin{pmatrix} \Sigma^+ \\ \Sigma^0 \\ \Sigma^- \end{pmatrix} \quad \begin{pmatrix} \pi^+ \\ \pi^0 \\ \pi^- \end{pmatrix}$$

$$\begin{aligned} 4\pi^- + p \rightarrow K^0 + \Sigma^0 : \\ |1, -1\rangle \left| \frac{1}{2}, \frac{1}{2} \right\rangle \rightarrow \left| \frac{1}{2}, -\frac{1}{2} \right\rangle |1, 0\rangle \\ \Leftrightarrow \sqrt{\frac{1}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle - \sqrt{\frac{2}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \rightarrow \sqrt{\frac{2}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle + \sqrt{\frac{1}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \end{aligned}$$

$$\begin{aligned} M = \langle I, M | T | I', M' \rangle \\ M_1 = \langle \pi^- p | T | K^0 \Sigma^0 \rangle \\ = \left(\sqrt{\frac{1}{3}} \left\langle \frac{3}{2}, -\frac{1}{2} \right| - \sqrt{\frac{2}{3}} \left\langle \frac{1}{2}, -\frac{1}{2} \right| \right) T \left(\sqrt{\frac{2}{3}} \left| \frac{3}{2}, -\frac{1}{2} \right\rangle + \sqrt{\frac{1}{3}} \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \right) \\ = \frac{\sqrt{2}}{3} T_{3/2} - \frac{\sqrt{2}}{3} T_{1/2} \\ M_2 = \langle \pi^- p | T | K^+ \Sigma^- \rangle = \frac{1}{3} T_{3/2} + \frac{2}{3} T_{1/2} \\ M_3 = \langle \pi^+ p | T | K^+ \Sigma^+ \rangle = T_{3/2} \end{aligned}$$

$$\sigma \propto M^2$$

$$\begin{aligned} I = \frac{3}{2} \Rightarrow T_{1/2} = 0 \Rightarrow \frac{2}{9} : \frac{1}{9} : 1 \Leftrightarrow 2 : 1 : 9 \\ I = \frac{1}{2} \Rightarrow T_{3/2} = 0 \Rightarrow \frac{2}{9} : \frac{4}{9} : 0 \Leftrightarrow 1 : 2 : 0 \end{aligned}$$