

Hadron and Nuclear Physics

Exercise sheet 2

Winter term 2013/14

Hand-in: Monday, Nov 18th 2013, Room 108

Problem 2.1: Magnetic Dipole Moments of Nucleons (9 points)

(a) Calculate the magnetic dipole moments of protons and neutrons considering the following information:

- The three constituent quarks carry one third of the entire nucleon mass.
- Quarks are pointlike fermions whose magnetic dipole moment can be calculated in analogy to the electron from their spin, charge and mass (Dirac particles).
- The magnetic moment of a three-quark state is obtained from the following quantum-mechanical relation:

$$\mu = 2\langle q_1 q_2 q_3 | \mu_1 S_{z1} + \mu_2 S_{z2} + \mu_3 S_{z3} | q_1 q_2 q_3 \rangle = 2 \sum_{i=1}^3 \langle q_i | \mu_i S_z | q_i \rangle$$

- The wave function of a nucleon with spin \uparrow is given by:

$$|N_{\uparrow}\rangle = \sqrt{2/9} (|1_{\uparrow}2_{\uparrow}3_{\downarrow}\rangle + |3_{\downarrow}1_{\uparrow}2_{\uparrow}\rangle + |2_{\uparrow}3_{\downarrow}1_{\uparrow}\rangle) \\ - \sqrt{1/18} (|1_{\downarrow}2_{\uparrow}3_{\uparrow}\rangle + |3_{\uparrow}1_{\downarrow}2_{\uparrow}\rangle + |2_{\uparrow}3_{\uparrow}1_{\downarrow}\rangle + |1_{\uparrow}2_{\downarrow}3_{\uparrow}\rangle + |3_{\uparrow}1_{\uparrow}2_{\downarrow}\rangle + |2_{\downarrow}3_{\uparrow}1_{\uparrow}\rangle)$$

(where for the proton $|123\rangle \equiv |ud\rangle$, for the neutron $|123\rangle \equiv |ddu\rangle$)

Compare the magnetic moments and their ratio with the experimental values for neutron and proton.

Problem 2.2: Exotic States (9 points)

(a) Which of these objects can exist? Why?

$$\bar{b}c, \quad uug, \quad \bar{c}cg, \quad gg, \quad ggg, \quad \bar{u}\bar{u}dd, \quad uudd, \quad uudd\bar{s}, \quad uuddss$$

(b) In which kind of reactions do you expect to find gluons and hybrids? How can you distinguish between a hybrid and a meson? Why is this so difficult?

Problem 2.3: Missing Mass (9 points)

(a) In an experiment protons with a kinetic energy of 300 MeV are fired at a hydrogen target at rest. The outgoing protons are detected in two detectors, which are arranged symmetrically around the beam at an angle of 12° . Pairs of protons with the energies, which you can find in the table, are measured. Which reaction do occur, elastic scattering, photon production or production of a particle (if so, which particle?)?

T_3/MeV	80.9	70	50.2	64.2	78.4	38.5	60.7	75.5	29.7	59.7	73.7	25.9	23.7
T_4/MeV	80.9	70	50.2	77.1	83.3	64.6	84.5	85.7	80.5	88.1	87.3	90.1	96.4

Problem 2.4: Charmonium (9 points)

- (a) Draw the level scheme of charmonium. Which quantum numbers do the different states have?
- (b) Compared to the energy levels of positronium, the scheme of charmonium is very similar for the low lying states ($n = 1, 2$), whereas for higher lying states the behaviour of positronium and charmonium is very different. What can one conclude from this observation?

- (c) Why are the partial widths of the ϕ ($\Gamma(\phi \rightarrow 3\pi) \approx 650 \text{ keV}$) and the J/ψ ($\Gamma(J/\psi \rightarrow \text{hadrons}) \approx 82 \text{ keV}$) so small compared to the width of the ρ ($\Gamma(\rho \rightarrow \pi\pi) \approx 150 \text{ MeV}$)? Draw the decays schematically with quark and gluon lines.
- (d) In the BELLE-Collaboration, the reaction $e^+e^- \rightarrow J/\psi X$ was studied. Here, the J/ψ was selected with a momentum of $|\vec{p}_{\text{CMS}}| > 2 \text{ GeV}$.




Figure 1: Mass distribution $M_{\text{Recoil}}(J/\psi) = \sqrt{(E_{\text{CMS}} - E_{J/\psi})^2 - \vec{p}_{J/\psi}^2}$ in the reaction $e^+e^- \rightarrow J/\psi X$ (K. Abe et al., BELLE-Collaboration, arXiv:hep-ex/0507019).

- Why did one use this momentum cut?
- Which quark content does the X have?
- Interpret the mass distribution in the picture. Which particles can you see there? Which spin, relative orbital and total angular momentum do they have?

2 Übung: Besprechung

2.1 Aufgabe

⇒ Griffith

$$|p\rangle = \underbrace{\frac{2}{3\sqrt{2}}|u\uparrow u\uparrow d\downarrow\rangle}_a - \underbrace{\frac{1}{3\sqrt{2}}|u\uparrow u\downarrow d\uparrow\rangle}_b - \underbrace{\frac{1}{3\sqrt{2}}|u\downarrow u\uparrow d\uparrow\rangle}_c + \underbrace{\text{permutation}}_{\Rightarrow 3*(a+b+c)}$$

$$\hat{S}_z|q_\uparrow\rangle = \hat{S}_z|q_i\rangle \underbrace{\frac{1}{2}}_S, \underbrace{\frac{1}{2}}_{m_s} \rangle = \frac{\hbar}{2}|q_p\rangle$$

$$\mu_A = \left(\frac{2}{3\sqrt{2}}\right)^2 \cdot 2 \sum_i \langle u\uparrow u\uparrow d\downarrow | \mu_i \hat{S}_z | u\uparrow u\uparrow d\downarrow \rangle = \frac{2}{9} \cdot 2 \left(\frac{\mu_u}{2} + \frac{\mu_u}{2} - \frac{\mu_d}{2} \right)$$

$$\mu_B = \left(\frac{1}{3\sqrt{2}}\right)^2 \cdot 2 \sum_i \langle u\uparrow u\downarrow d\uparrow | \mu_i \hat{S}_z | u\uparrow u\downarrow d\uparrow \rangle = \frac{1}{18} \cdot 2 \left(\frac{\mu_u}{2} - \frac{\mu_u}{2} + \frac{\mu_d}{2} \right) = \frac{1}{18} \mu_d$$

$$\Rightarrow \mu_C = \mu_B = \frac{1}{18} \mu_d$$

$$\mu_p = 3 \left(\frac{2}{9} (2\mu_u - \mu_d) \right) + \frac{1}{18} \mu_d + \frac{1}{18} \mu_d = \frac{1}{3} (4\mu_u - \mu_d) \Rightarrow \mu_d = \frac{1}{3} (4\mu_d - \mu_u)$$

$$\text{Vereinfachungen: } m_u = m_d = m_q = \frac{1}{3} m_N, \text{ Dirac-Teilchen: } \mu = \frac{Q\hbar}{2mc}$$

$$\text{Kernmagneton (Wenn Photon Dirac-Teilchen): } \mu_N = \frac{e\hbar}{2m_p c}$$

$$\Rightarrow \mu_u = \frac{2}{3} \frac{e\hbar}{\frac{1}{3} m_p c} = 2\mu_N, \quad \mu_d = -\frac{1}{3} \frac{e\hbar}{\frac{1}{3} m_p c} = -\mu_N$$

$$\mu_p = \frac{1}{3} (4\mu_u - \mu_d) = 3\mu_N \quad (2.793\mu_N), \quad \mu_n = \frac{1}{3} (4\mu_d - \mu_u) = -2\mu_N \quad (-1.913\mu_N)$$

$$\frac{\mu_u}{\mu_d} = -\frac{2}{3} \quad (0.68)$$

2.2 Aufgabe

a) Farbladung! (e.g. uug verboten, da u je 1 Farbe, g jedoch Farbe + Anti-farbe)

b) Was ist uuddss? $p\bar{\Xi}^-, n\bar{\Xi}^0, \Lambda\Lambda, \Sigma^+\Sigma^-$ (DiBaryon)

Beispiel: $\pi^0 \rightarrow \gamma\gamma$: $J^{PC} : 0^{-+} \rightarrow (1^{--})^2$ (nicht $\gamma\gamma$!)

$\rho \rightarrow \pi^+\pi^-$: $P(\rho^0) = \eta_{\pi^+}\eta_{\pi^-} \cdot (-1)^l$

$P(\rho) = -1 \Rightarrow l = 1, 3, \dots$, $P(\rho) = 1 \Rightarrow l = 2, 4, \dots$

Glueball:

zerfällt nur stark \Rightarrow schnell \Rightarrow Breite \Rightarrow Andere Kanäle überlagern

Aber: Flavorblind!

Bsp: $m(G) = 20\text{GeV}$

$$Br(G \rightarrow B^0 \bar{B}^0) \approx Br(G \rightarrow D^0 \bar{D}^0) \approx Br(G \rightarrow K^0 \bar{K}^0)$$

2.3 Aufgabe

$$\text{Rutherford: } \sigma(\vartheta) = \left(\frac{Q_1 Q_2}{4T} \right)^2 \frac{1}{\sin^4(\frac{\vartheta}{2})}$$

$$\text{Fällt sehr schnell ab: } \frac{\sigma(12^\circ)}{\sigma(13^\circ)} = 5 \cdot 10^{-5}$$

$$\text{Missing Mass! Energieerhaltung: } T + 2m_p = T_3 + T_4 + 2m_p + E_x \Rightarrow E_x = T - T_3 - T_4$$

Nun 4-Impuls-Erhaltung:


$$p_3 = \sqrt{(T_3 + m_3)^2 - m_p^2}, \quad p_4 = \sqrt{(T_4 + m_4)^2 - m_p^2}$$

$$\text{Transversal: } p_T = 0 = p_{T,3} + p_{T,4} + p_{T,X} \Rightarrow p_{T,X} = -p_{T,3} - p_{T,4}$$

$$\text{Longitudinal: } p_Z = p_{Z,3} + p_{Z,4} + p_{Z,X} \Rightarrow p_{Z,X} = p_Z - p_{Z,3} - p_{Z,4}$$

$$p_{T,3} = \sin(\vartheta)p_3, \quad p_{Z,3} = \cos(\vartheta)p_3, \quad p_X^2 = p_{T,X}^2 + p_{Z,X}^2 \Rightarrow M_x = \sqrt{E_x^2 - p_x^2}$$

$$M_x \approx m_\pi \Rightarrow pp \rightarrow pp\pi^0, \quad M_x \approx 0 \Rightarrow pp \rightarrow pp\gamma$$



2.4 Aufgabe

$$e^+e^- \rightarrow Y(4s) \rightarrow J/\psi X$$

$$X = \pi^+\pi^-, \pi^0\pi^0, \eta_c, \eta_c\pi^+\pi^-, \dots (J/\psi \rightarrow \mu^+\mu^- : Br \approx 10^{-4})$$

$$\text{CMS: } \begin{pmatrix} E_{CMS} \\ 0 \end{pmatrix} = \begin{pmatrix} E_\psi \\ \vec{p}_\psi \end{pmatrix} + \begin{pmatrix} E_x \\ \vec{p}_x \end{pmatrix} \Rightarrow (E_{CMS} - E_\psi)^2 - \vec{p}_\psi^2 = M_x^2$$