II. Physikalisches Institut

JUSTUS-LIEBIG-UNIVERSITÄT GIESSEN

IONAS

GSI

HELMHOLTZ GEMEINSCHAFT

Cancer therapy with photons, protons and heavy ions

Julian Bergmann
Justus-Liebig-University Gießen

Overview

- Introduction
- Overview
- Particles in radiation therapy
- Raster scan method
- Summary

Introduction

Motivation

→ tumor therapy requires

→ affected tissues to be removed

→ spreading to get stopped

→ not killing the patient while doing that.

→ chirurgic treatment needs accessibility to tumor

→ chemical treatment stresses the overal body much

→ radiation therapy can target more precisely

History

1895 Wilhelm Conrad Röntgen „a new kind of radiation“

1897 first medical treatment with X-rays

1957, Berkeley, first proton beam therapy

1975, Berkeley, first use of heavy ions.

1979, GSI Darmstadt, SIS12 (SIS18 in 1984) for 1.4 GeV/n

1993, Chiba, carbon beam therapy department founded

1993-97, GSI Darmstadt, carbon beam therapy facility founded

→ development of raster scan procedure

→ development of PET quality control

→ measurements of ion species dependand RBE

2002, Hyogo, carbon beam facility founded

2009, Heidelberg, HIT carbon beam facility, ~ 500 patients per year

2012, Kiel, and 2015 in Marburg, carbon beam facility founded

Overview

Ionizing particles hit Cell

→ Low energy transfer creates oxygen radicals

→ DNA gets damaged beyond repair

→ Cell division is disturbed (mitotic cell death)

→ Cell dies (Apoptosis, blebbing, nucleus fragmentation)

Effectiveness depends on:

→ deposited energy

→ Choice of particles

→ technical application

→ biological reaction to particle

Particle in Radiation Therapy

Particles used for radiation therapy:


→ Photons

→ Electrons

→ Neutrons


→ Protons

→ Ions



https://de.wikipedia.org/wiki/Tiefendosiskurve

Particle Interaction: Photons



Photons (<23MeV)

→ Easy to produce

→ Xrays (~120 keV) mostly photoeffect

→ energy deposition exponentially decreasing

→ additional energy deposition by secondary electrons

→ healthy tissue before and after target tissue is affected

Photoeffect: bound electron absorbs photon and gets released

$$E_{kin} = h \cdot f - E_{\text{Bindung}}$$

Compton: photon scatters at free electron and looses energy

$$\Delta\lambda = \frac{h}{m_e c}\left(1 - \cos\phi\right) = \lambda_C\left(1 - \cos\phi\right) .$$

Pair production: if $E_\gamma > 2m_e c^2$, a electron positron pair can be produced inside the field of a nucleus

Particle Interaction: Electrons

Electrons


→ high energy: Bremsstrahlung

→ low energy: ionisation and excitation.

→ maximum range (~ 0.5 cm/MeV)

→ less energy deposition per collision than photons

→ elastic collision lead to additional beam spread



http://pdg.lbl.gov/2013/reviews/rpp2013-rev-passage-particles-matter.pdf

Particle Interaction: Protons

Protons


→ produced by particle accelerator

→ high energy: inelastic collisions with target electrons

→ lower energy: inelastic collisions with nuclei

→ elastic scattering at nuclei: beam spread

→ energy deposition decreases with square of velocity



https://de.wikipedia.org/wiki/Tiefendosiskurve

→ result: proton slows down until stop at energy deposition peak

→ „Bragg peak“

$$\frac{1}{\rho}\frac{dE}{dx} = Kz^2\frac{Z}{A}\frac{1}{\beta^2}\left[\frac{1}{2}\ln\frac{2m_e c^2\beta^2\gamma^2 T_{\max}}{I^2} - \beta^2 - \frac{\delta(\beta\gamma)}{2}\right]$$

→ low dose until maximum bragg peak, afterwards 0 dosage

→ bragg peak position depending on proton energy

→ less lateral scattering than X-Rays or electrons

Particle Interaction: Ions

Ions

→ similar processes as protons

→ higher mass

→ less inelastic scattering per dx

→ less dE/dx until bragg peak

→ little lateral scatter effect

→ higher linear energy transfer (LET) at bragg peak

→ much narrower bragg peak

→ also direkt cell damage effect by nuclear collision


→ nuclear reactions with tissue: dE/dx > 0 behind bragg peak

→ higher energy needed for same bragg peak distance

→ common: carbon ions

→ highest overlap between biological effectiveness and energy deposition

Particle Interaction: Comparison



History of the heavy ion therapy at GSI. Kraft G

Particle Interaction: Neutrons

Neutrons

→ need nuclear reaction to produce (reactors, cyclotrons)

→ collisions at low energy produce protons in tumor cell

→ better for low oxygen cells

→ only 1/3 effective dose of protons needed

→ similar deposition curve as X-Rays


→ using Bor in cell can be more effective:

$$n + {}^{10}\mathrm{B} \longrightarrow {}^{7}\mathrm{Li} + \alpha$$

→ cell gets destroyed directly

→ most dE is at surface

→ high biological effectiveness and bad in depth dose distribution means high damage at healthy cells




http://rpd.oxfordjournals.org/content/137/1-2/149/F1.large.jpg

Raster Scan Method: Idea

Idea:

→ dipolmagnets scan tissue lateral

→ protons/ions: energy determines depth (bragg peak)



Tumortherapie mit Ionenstrahlen, GSI, Kraft G

Problem:

→ energy deposition spread in tissue

→ minimize energy deposition at healthy tissues

→ tissue species reacts differently to same dosage

→ hard to target moving body parts (e.g. lung, intestines)

Raster Scan Method: RBE

RBE: Relative Biological Effectivenes

→ quotient of dose to a reference type of radiation with same biological effect.


→ increases with LET (DNA double-strand breaks)

→ highly depends on ion species and type of tissue


→ biological effective dose = RBE * energy deposition

→ determined experimental

→ carbon has RBE maximum at bragg peak



History of the heavy ion therapy at GSI. Kraft G, courtesy Michael Kraemer, GSI



Wilma K. Ewyrather, Gerhard Kraft, GSI Darmstadt – Abt. Biophysik

Raster Scan Method: Spread Out Bragg Peak


Spread out bragg peak:

→ overlaying ion beams with different energies

→ spreading bragg peak longitudinal

→ keeping target-dose constant

→ minimizing collateral dose



http://www.nature.com/bjc/journal/v93/n8/images/6602754f1.jpg

Raster Scan Method: Application

Raster Scan procedure

→ tumor division into slices

~ 100 slices

→ each slice is divided into raster cells

~ 10 - 30000 raster cells per slice

→ beam diameter ~ 3 raster cells

→ from RBE & energy deposition,


to be applied energy is computed

Example:

→ complete application time: 5-10 min

→ daily application for 20 days

→ beam size ~ 6 mm



Tumortherapie mit schweren Ionen, GSI, Gerhard Kraft

Raster Scan method: PET

PET (Positron Emission Tomography) Verification:


→ quality control: which dosage was applied where

→ nuclear collisions produce instabile atoms.


→ ^{10}C, ^{11}C and ^{15}O decay emitting positrons

→ positrons decay into 2 gamma ~ 511 keV

→ detection gives estimate of beam particle reach



Facilities



http://www.hirt-japan.info/en/what/japan.html

→ during 2013 105.000 patients successfully treated with protons

→ 13.000 patients per year treated by heavy ions

→ ~ 49 working proton beam facilities

→ ~ 10 working ion beam facilities using carbon

Summary

→ radiation therapy is nowadays established cancer therapy method

→ photons, electrons, neutrons, protons and ions

→ heavy ion beams promise to be most effective method (situational)

→ RBE, raster scan method and PET further increases precision

→ facilities for heavy ion beam production are still expensive,

but centers are build all over the world

Papers and references

History of the heavy ion therapy at GSI. Kraft G. https://three.jsc.nasa.gov/articles/Krafts_GSI.pdf. Date posted: 04-26-2013.

Forschungsreaktor FRM II: Krebsbehandlung mit Neutronen fragwürdig, Karin Wurzbacher, 2003, Umweltinstitut München

Wechselwirkung und Reichweite von Strahlung, Hartmut Zabel, 2003, Presentation

Physik der Partikeltherapie, Petra Knappe-Kagan, 2012, Institut für Medizinische Physik Klinikum Nürnberg, Presentation

Tumortherapie mit Ionenstrahlen, Gerhard Kraft, 2005, Biophysik GSI Darmstadt

Particle Beam Cancer Therapy: The Promise and Challenges, 2014, Karen McNulty Walsh, article on RHIC website: https://www.bnl.gov/rhic/news2/news.asp?a=4672&t=today

Passage of particles through matter, 2013, . Beringer et al. (PDG), http://pdg.lbl.gov/2013/reviews/rpp2013-rev-passage-particles-matter.pdf